

Different versions of mapping class groups of surfaces

Søren K. Boldsen

August 16, 2009

1 Introduction

Let F be a compact connected smooth surface, possibly with boundary and not necessarily oriented. The objects of study in this paper are

$$\begin{aligned}\mathrm{Diff}(F, \{\partial F\}) &= \{\varphi : (F, \partial F) \longrightarrow (F, \partial F) \mid \varphi \text{ is a diffeomorphism}\}, \\ \mathrm{Top}(F, \{\partial F\}) &= \{\varphi : (F, \partial F) \longrightarrow (F, \partial F) \mid \varphi \text{ is a homeomorphism}\}, \\ \mathrm{hAut}(F, \{\partial F\}) &= \{\varphi : (F, \partial F) \longrightarrow (F, \partial F) \mid \varphi \text{ is a homotopy equivalence}\}.\end{aligned}$$

The main theorem of this paper is the following:

Theorem 1.1. *Let F be a compact surface and not a sphere, a disk, a cylinder, a Möbius band, a torus, a Klein bottle, or $\mathbb{R}P^2$. Then*

$$\pi_0(\mathrm{Diff}(F, \{\partial F\})) \xrightarrow{\cong} \pi_0(\mathrm{Top}(F, \{\partial F\})) \xrightarrow{\cong} \pi_0(\mathrm{hAut}(F, \{\partial F\}))$$

are bijections.

This result is far from new, but this paper will present a thorough and self-contained proof of the following bijection

$$\pi_0(\mathrm{Diff}(F, \{\partial F\})) \xrightarrow{\cong} \pi_0(\mathrm{hAut}(F, \{\partial F\})). \quad (1)$$

To get the Main Theorem from this result, we will use the result of [Epstein] Thm 6.4 without proof.

We consider slightly different versions of the groups, where we assume F is oriented in the last two groups:

$$\begin{aligned}\mathrm{Diff}(F, \partial F) &= \{\varphi \in \mathrm{Diff}(F, \{\partial F\}) \mid \varphi|_{\partial F} = \mathrm{id}\}, \\ \mathrm{Diff}_+(F, \{\partial F\}) &= \{\varphi \in \mathrm{Diff}(F, \{\partial F\}) \mid \varphi \text{ is orientation-preserving}\}, \\ \mathrm{Diff}_+(F, \partial F) &= \mathrm{Diff}(F, \partial F) \cap \mathrm{Diff}_+(F, \{\partial F\}),\end{aligned}$$

and similar for Top and hAut . By orientation-preserving we mean that the orientation class $[F, \partial F] \in H_2(F, \partial F)$ is preserved by φ_* . From the Main Theorem we easily deduce

Theorem 1.2. *Let F be a compact surface and not a sphere, a disk, a cylinder, a Möbius band, a torus, a Klein bottle, or $\mathbb{R}P^2$. Then there are bijections*

- (1) $\pi_0(\text{Diff}(F, \partial F)) \xrightarrow{\cong} \pi_0(\text{Top}(F, \partial F)) \xrightarrow{\cong} \pi_0(\text{hAut}(F, \partial F)),$
- (2) $\pi_0(\text{Diff}_+(F, \{\partial F\})) \xrightarrow{\cong} \pi_0(\text{Top}_+(F, \{\partial F\})) \xrightarrow{\cong} \pi_0(\text{hAut}_+(F, \{\partial F\})),$
- (3) $\pi_0(\text{Diff}_+(F, \partial F)) \xrightarrow{\cong} \pi_0(\text{Top}_+(F, \partial F)) \xrightarrow{\cong} \pi_0(\text{hAut}_+(F, \partial F)).$

The standard definition of the mapping class group of a surface F is $\Gamma(F) = \pi_0(\text{Diff}_+(F, \partial F))$. The last part of Theorem 1.2 shows that it does not matter whether one considers diffeomorphisms, homeomorphisms, or even homotopy equivalences, when working in the mapping class group.

It is a pleasure to thank Jørgen Tornehave for many fruitful discussions and help during my work on this paper.

Contents

1	Introduction	1
2	Preliminaries	3
3	Surjectivity	5
4	Injectivity	11
5	Proof of the Main Theorem	16

2 Preliminaries

Definition 2.1. An isotopy ψ of F is a path in $\text{Top}(F, \{\partial F\})$, i.e. $\psi : F \times I \longrightarrow F$ is continuous map such that $\psi_t = \psi(-, t) : F \longrightarrow F$ is a homeomorphism for all $t \in I$, and we say that ψ_0 and ψ_1 are isotopic.

An isotopy is smooth if we can exchange homeomorphism with diffeomorphism in the above. We then say that ψ_0 and ψ_1 are smoothly isotopic.

Lemma 2.2. *Let $f : S^1 \longrightarrow S^1$ an orientation preserving diffeomorphism. Then f is smoothly isotopic to the identity via a smooth isotopy $f_t : S^1 \times I \longrightarrow S^1$ such that the function $F : S^1 \times I \longrightarrow S^1 \times I$ given by $F(z, t) = (f_t(z), t)$ is a diffeomorphism, and*

$$f_t(z) = \begin{cases} f(z), & \text{for } 0 \leq t < \varepsilon, \\ z, & \text{for } 1 - \varepsilon < t \leq 1. \end{cases}$$

Proof. Since f is smooth it defines a smooth function $\tilde{f} : \mathbb{R} \longrightarrow \mathbb{R}$ by lifting f under the universal covering $\exp : \mathbb{R} \longrightarrow S^1$. Now take a smooth bump function $\rho : I \longrightarrow I$ satisfying

$$\rho(t) = \begin{cases} 0, & t \leq \varepsilon, \\ 1, & t \geq 1 - \varepsilon. \end{cases}$$

Let $\tilde{F} : \mathbb{R} \times I \longrightarrow \mathbb{R}$ be given by $\tilde{F}(x, t) = \rho(t)\tilde{f}(x) + (1 - \rho(t))x$. This now defines an isotopy from $\tilde{f}$ to the identity, and $F(\exp(x), t) = (\exp(\tilde{F}(x, t)), t)$ is a diffeomorphism. $\square$

The idea of the following proof is due to J. Alexander.

Lemma 2.3. *Let D be a disk and N a collar neighborhood of the boundary. Suppose $f : D \longrightarrow D$ is a homotopy equivalence which restricts to an orientation preserving diffeomorphism of N of the form $f(z, t) = (f(z), t)$ for $(z, t) \in N$. Then f is homotopic to a diffeomorphism relative to a smaller collar neighborhood.*

Proof. We can assume $f : D \longrightarrow D$, where $D = \{z \in \mathbb{R}^2 \mid |z| \leq 1 + \varepsilon\}$, and $N = \{z \in D \mid 1 - \varepsilon < |z| \leq 1 + \varepsilon\}$. The tubular coordinates on N are $s \in [0, 2\pi]$ and $t \in (-\varepsilon, \varepsilon]$. We first construct a homotopy φ_x , $x \in [0, 1]$, which is constant in x outside N , from f to a function g such that $g(s, 0) = (s, 0)$ in tubular coordinates. We use the isotopy $f_x(s)$ from Lemma 2.2, and set

$$\varphi_x(s, t) = (f_{x(1 - \frac{1}{\varepsilon}|t|)}(s), t), \quad t \in (-\varepsilon, \varepsilon]$$

in tubular coordinates. Then $\varphi_0 = f$ and $\varphi_1(s, 0) = (s, 0)$, and φ_x is the identity on a collar neighborhood of ∂D by Lemma 2.2.

We now make a homotopy ψ_x , $x \in [0, 1]$, from g to the function h satisfying

$$h(z) = \begin{cases} g(z), & |z| > 1; \\ z, & |z| \leq 1. \end{cases}$$

Let $D' = \{z \in D \mid |z| \leq 1\}$, and define the solid cone

$$C = \{(z, x) \subseteq D \times I \mid |z| \leq 1 - x\}$$

with bottom $D' \times \{0\}$ and top $(0, 1)$, and set

$$\psi_x(z) = \begin{cases} (1-x)f(\frac{z}{1-x}), & (z, x) \in C, \\ z, & (z, x) \in (D' \times I) \setminus C, \\ g(z), & (z, x) \in (D \setminus D') \times I. \end{cases}$$

This is clearly continuous and constitutes a homotopy from g to h through maps which are the identity on a collar neighborhood of ∂D , since g is. We claim $h : D \rightarrow D$ is a diffeomorphism. Clearly, $h : D' \rightarrow D'$ is a diffeomorphism, and by Lemma 2.2, h is smooth on D , and for $|z| > 1$, $h = g$ is a diffeomorphism $D \setminus D' \rightarrow D \setminus D'$. $\square$

A result we will use repeatedly is the following smooth version of the Schönflies curve theorem.

Lemma 2.4. *Let $f : S^1 \rightarrow F$ be a smoothly embedded simple closed curve homotopic to zero in a surface F . Then the closure of the interior of $f(S^1)$ is a smoothly embedded disk in F .*

Proof. By Thm 1.7 in [Epstein] we know that f separates F into two components, and that one of them (call it D') is homeomorphic to a disk D^2 . Thus D' is a connected orientated smooth 2-manifold with 1 boundary component and with Euler characteristic $\chi(D') = 1$. Now by the classification of smooth surfaces, D' is a smooth disk. $\square$

Definition 2.5. Let α be a smoothly embedded 1-submanifold in a surface F . By the surface cut up along α , denoted $F \setminus \alpha$, we will mean the surface with boundary $F \setminus N(\alpha)$, where $N(\alpha)$ is a tubular neighborhood of α in F .

Lemma 2.6. *Let $\alpha : (I, \partial I) \rightarrow (F, \partial F)$ be a simple curve in a surface F . If the cut-up surface $F \setminus \alpha(I)$ is disconnected, then the induced map $\alpha_* : H_1(I, \partial I) \rightarrow H_1(F, \partial F)$ is the zero map.*

Proof. Let $\bar{\alpha} = \alpha(I) \subseteq F$, and consider the long exact sequence for the triple $(\partial F, \bar{\alpha} \cup \partial F, F)$:

$$H_1(\bar{\alpha} \cup \partial F, \partial F) \xrightarrow{i_*} H_1(F, \partial F) \xrightarrow{j_*} H_1(F, \bar{\alpha} \cup \partial F) \longrightarrow H_0(\bar{\alpha} \cup \partial F, \partial F)$$

Here $H_0(\alpha \cup \partial F, \partial F) = 0$, so j_* is surjective. Also $H_1(F, \partial F) \cong \mathbb{Z}^{2g+r-1}$ for $F = F_{g,r}$. Since $F \setminus \bar{\alpha}$ is not connected, we can write $F \setminus \bar{\alpha} = F_1 \sqcup F_2$, and by excision,

$$\begin{aligned} H_1(F, \alpha \cup \partial F) &\cong H_1(F_1 \sqcup F_2, \partial F_1 \sqcup \partial F_2) \cong H_1(F_1, \partial F_1) \oplus H_1(F_2, \partial F_2) \\ &\cong \mathbb{Z}^{2g_1+r_1-1} \oplus \mathbb{Z}^{2g_2+r_2-1}. \end{aligned}$$

Here $g = g_1 + g_2$ and $r + 1 = r_1 + r_2$, so since j_* is surjective, we conclude that j_* is an isomorphism. Thus $i_* = 0$, and the following diagram shows that $\alpha_* = 0$:

$$\begin{array}{ccc} H_1(I, \partial I) & \xrightarrow{\alpha_*} & H_1(F, \partial F) \\ \alpha_* \downarrow & \nearrow i_* & \\ H_1(\bar{\alpha} \cup \partial F, \partial F) & & \end{array}$$

□

3 Surjectivity

In this section we will prove that the map in (1) is surjective, i.e. a homotopy equivalence of a surface F is homotopic to a diffeomorphism. We first prove this for surfaces with non-empty boundary, and then use this to obtain the proof for closed surfaces. The result for surfaces with non-empty boundary is strongly inspired by [Hempel].

Theorem 3.1. *Let F and G be compact surfaces with non-empty boundaries. Suppose $\pi_1(F)$ is non-trivial. Let $f : (F, \partial F) \rightarrow (G, \partial G)$ be a map such that $f_* : \pi_1(F) \rightarrow \pi_1(G)$ is injective and $f|_{\partial F} : \partial F \rightarrow \partial G$ is a smooth embedding. Then there is a homotopy $f_t : (F, \partial F) \rightarrow (G, \partial G)$ with $f_0 = f$ and $f_1 : F \rightarrow G$ a diffeomorphism.*

Proof. First consider each boundary component J of F , and K of G where $f(J) \subseteq K$. We can assume each J and K has a collar neighborhood of the form $J \times [0, \varepsilon]$ and $K \times [0, \varepsilon]$, where the map f has the form $f(x, t) = (f|_J(x), t)$, by gluing on small cylinders, extending f as desired, and smoothing out. Since f is continuous, it is homotopic to a smooth map, and we can choose the homotopy to be constant on the collar neighborhoods, so we can assume that f is smooth an embedding on a neighborhood of ∂F .

We are going to cut up G by a non-separating arc α (i.e. an embedded connected 1-manifold with boundary) connecting two boundary components of G in the image of f . We would like to cut up F by $f^{-1}(\alpha)$. To do this we must ensure that $f^{-1}(\alpha)$ is also an embedded 1-manifold. This holds if f is

transverse to α . By Thom's transversality theorem, f can be approximated by a smooth map g transverse to α arbitrarily close to f . Even better, g can be chosen such that $g|_A = f|_A$ for a closed subset $A \subseteq F$ on which the transversality condition on f is already satisfied. If we choose the arc α to have the form $\alpha = (x_0, t)$, $t \in [0, \varepsilon]$ on the collars $K \times [0, \varepsilon]$ for some $x_0 \in K$, then clearly we can take $A = \bigcup_{J \in \pi_0(F)} J \times [0, \varepsilon]$ in the above. Since the transverse map g is arbitrarily close to f , they are homotopic, and we can assume f is transverse to α .

Since $f|_{\partial F} : \partial F \rightarrow \partial G$ is an embedding we can see that $f^{-1}(\alpha)$ must consist of one arc in F and possibly a number of embedded circles, and as F is compact, there is a finite number of circles. Since f_* is injective, the circles must be null-homotopic in F , thus they must each bound a disk D_0 in F . Taking a slightly larger disk $D \supseteq D_0$, then $f(\partial D)$ must be contained in a tubular neighborhood of α . Since ∂D is disjoint from $f^{-1}(\alpha)$, all of $f(\partial D)$ is to the same side of α in the tubular neighborhood.

Now D is a disk and $f(\partial D)$ is contained in a disk $E \subseteq G$ on one side of α in the tubular neighborhood. Thus we can make a map $h : D \rightarrow G$ with $h(D) \subseteq E$ and such that $f|_{\partial D} = h|_{\partial D}$. This gives a map $H : S^2 \rightarrow G$ by mapping the lower hemisphere by f and the upper hemisphere by h . Since G is not S^2 or $\mathbb{R}P^2$, we know $\pi_2(G) = 0$, so the map H can be extended to a map $D^3 \rightarrow G$, thus giving a homotopy from f to h . This will reduce the number of circles in the preimage, and we can thus assume that $f^{-1}(\alpha)$ is just an arc in F . By transversality we can assume that we have a tubular neighborhood of $f^{-1}(\alpha)$ mapping to a tubular neighborhood of α .

We can now cut F along $f^{-1}(\alpha)$ and G along α , to obtain $\hat{F}$ and $\hat{G}$. After cutting up F and G along an arc, we will actually have manifolds with corners, $\hat{F}$ and $\hat{G}$. But clearly we can smooth out these corners inside the collar neighborhoods where $f : \hat{F} \rightarrow \hat{G}$ is smooth.

Now we would like to show that the process will not separate F . Consider the situation when we cut up along a non-separating arc α in G . We can parametrise α and think of it as a function $\alpha : (I, \partial I) \rightarrow (G, \partial G)$. This induces a map $\alpha_* : H_1(I, \partial I; \mathbb{Z}_2) \rightarrow H_1(G, \partial G; \mathbb{Z}_2)$. The condition that α is nonseparating translates as $\alpha_* \neq 0$. By the above we can assume that $f^{-1}(\alpha)$ is a single arc, which we parametrize as $\tilde{\alpha} : (I, \partial I) \rightarrow (F, \partial F)$:

$$\begin{array}{ccc} & (I, \partial I) & \\ \tilde{\alpha} \swarrow & & \searrow \alpha \\ (F, \partial F) & \xrightarrow{f} & (G, \partial G) \end{array}$$

On homology this induces the commutative diagram

$$\begin{array}{ccc}
& H_1(I, \partial I; \mathbb{Z}_2) & \\
\tilde{\alpha}_* \swarrow & & \searrow \alpha_* \\
H_1(F, \partial F; \mathbb{Z}_2) & \xrightarrow{f_*} & H_1(G, \partial G; \mathbb{Z}_2)
\end{array}$$

But since $\alpha_* \neq 0$ we get $\tilde{\alpha}_* \neq 0$ and thus by Lemma 2.6, $\tilde{\alpha} \subseteq F$ is nonseparating.

Now we show that $f_* : \pi_1(\hat{F}) \rightarrow \pi_1(\hat{G})$ is still injective after cutting up. We use that F is homotopic to $\hat{F} \cup I$, where I is a small interval connecting two points $b_0, b_1 \in \partial \hat{F}$. Using that $\hat{F}$ is connected we choose a path J in $\hat{F}$ from b_0 to b_1 , such that $I \cup J$ form a loop. Now $F \simeq \hat{F} \vee S^1$ (by contracting J in $\hat{F}$ to a point). Then $i_* : \pi_1(\hat{F}) \rightarrow \pi_1(F)$ is injective, since $i_* : \pi_1(\hat{F}) \rightarrow \pi_1(F) = \pi_1(\hat{F}) * \mathbb{Z}$ is just the inclusion in the first factor by van Kampen's theorem. Now it follows from the commutative diagram

$$\begin{array}{ccc}
\pi_1(\hat{F}) & \xhookrightarrow{i_*} & \pi_1(F) \\
\downarrow \hat{f}_* & & \downarrow f_* \\
\pi_1(\hat{G}) & \xrightarrow{i_*} & \pi_1(G)
\end{array}$$

that $\hat{f}_* : \pi_1(\hat{F}) \rightarrow \pi_1(\hat{G})$ is also injective.

It remains to show that by cutting up F and G they have to become disks at the same time. Firstly if G is a disk, then $f_* : \pi_1(F) \rightarrow \{1\}$ is injective, so $\pi_1(F) = \{1\}$, and this implies that F is also a disk (since F is a surface with boundary). Conversely, if G is not a disk then neither is F , since given a non-separating arc α in G we have shown above that there exists a non-separating arc in F .

We are down to the case where f is a map from a disk to a disk that is smooth in a collar of the boundary, and this case is handled by Lemma 2.4. We can glue the resulting smooth maps on the pieces together again, since the collar neighborhoods of the boundary of each piece (where the map is smooth) are fixed by the homotopy from Lemma 2.4. So we are done. $\square$

Corollary 3.2. *Let F and G be compact surfaces with non-empty boundaries. Suppose $\pi_1(F)$ is non-trivial. Let $f : (F, \partial F) \rightarrow (G, \partial G)$ be a map such that $f_* : \pi_1(F) \rightarrow \pi_1(G)$ is injective and $f|_{N(\partial F)} : N(\partial F) \rightarrow N(\partial G)$ is a smooth embedding, where $N(-)$ denotes a neighborhood. Then there is a homotopy $f_t : (F, \partial F) \rightarrow (G, \partial G)$ with $f_0 = f$ and $f_1 : F \rightarrow G$ a diffeomorphism, such that $f_t = f_0$ on a neighborhood of ∂F .*

Proof. Use the proof above, but skip the first part which proves that $f|_{N(\partial F)} : N(\partial F) \longrightarrow N(\partial G)$ can be made into a smooth embedding. $\square$

Lemma 3.3. *Let $f_0, f_1 : S^1 \longrightarrow F$ be disjoint non-trivial two-sided embeddings in the surface F . Assume there exist $m, n \in \mathbb{Z}_+$ such that f_0^n and f_1^m represent the same free homotopy class in F . Then there is an embedding $\varphi : S^1 \times I \longrightarrow F$ such that $\varphi|_{S^1 \times \{i\}} = f_i$ for $i = 0, 1$, so f_0 and f_1 bound a cylinder.*

Proof. This is a special case of [Epstein], Lemma 2.4.

We start by cutting F up along f_0 and then gluing a disk onto each of the two new boundary components; let M be the connected component containing f_1 in the resulting surface. Since f_0 is null-homotopic in M , then so is f_0^n and thus f_1^m . Now we will show that f_1 is null-homotopic in M , so that it bounds a disk in M . First if $\partial M \neq \emptyset$, then $\pi_1(M)$ is a free group and thus if $f_1^m = 1$ then $f_1 = 1$. Else $\pi_1(M)$ is a free group modulo the relation $\partial = \prod_{i=1}^g [a_i, b_i] \in \pi_1(M)$ (oriented case) or $\partial = \prod_{i=1}^g a_i^2 \in \pi_1(M)$ (unoriented case). If $f_1^m = 1$ but $f_1 \neq 1$, $\pi_1(M)$ will have torsion, and by [Lyndon-Schupp] Prop. 5.18, the only case that allows for torsion is the unoriented case with $g = 1$. Then the component of M containing f_1 is an $\mathbb{R}P^2$, but then there are no non-trivial two-sided embeddings of S^1 . So there can be no torsion, and $f_1 = 1$ in $\pi_1(M)$.

The disk in M bounded by f_1 contains either one or two of the disks glued onto f_0 to form M , since f_1 was non-trivial in F . If the disk bounded by f_1 in M contains just one glued-on disk, then f_0 and f_1 together bound a disk blown up at one point; a cylinder in F . In particular, if f_0 is separating, then the disk bounded by f_1 in M contains just one glued-on disk, so we are done. Now if the disk bounded by f_1 in M contains two of the glued-on disks, then f_1 was separating in F , since we obtain F from M by removing the glued-on disks and gluing up along their boundaries. The cylinder can thus be obtained if we interchange f_0 and f_1 . $\square$

The condition in the preceding Theorem 3.1 about the map f being an embedding on the boundary is not essential if f is a homotopy equivalence, as we show next:

Lemma 3.4. *Suppose $f : (F, \partial F) \longrightarrow (G, \partial G)$ induces an isomorphism $f_* : \pi_1(F) \longrightarrow \pi_1(G)$, and suppose F is compact with $\partial F \neq \emptyset$ and is neither a disk, a cylinder nor a Möbius band. Then the following holds:*

- (i) *For all boundary components $J \subseteq \partial F$ and $K \subseteq \partial G$ such that $f(J) \subseteq K$, the composite $\mathbb{Z} \cong \pi_1(J) \xrightarrow{f} \pi_1(K) \cong \mathbb{Z}$ is multiplication by ± 1 ,*

and no two different boundary components in F are taken to the same boundary component in G .

- (ii) f is homotopic to a map $g : (F, \partial F) \longrightarrow (G, \partial G)$ with $g|_{\partial F} : \partial F \longrightarrow \partial G$ an embedding.

Proof. Let $J \subseteq F$ be a boundary component, and let $K \subseteq G$ be the boundary component with $f(J) \subseteq K$. We have a commutative diagram,

$$\begin{array}{ccc} \pi_1(J) & \xrightarrow{(f|_J)_*} & \pi_1(K) \\ \downarrow & & \downarrow \\ \pi_1(F) & \xrightarrow[f_*]{\cong} & \pi_1(G) \end{array} \quad (2)$$

Here, the vertical map $\pi_1(J) \longrightarrow \pi_1(F)$ is injective, since it is a non-zero map (as F is not a disk) from $\pi_1(J) \cong \mathbb{Z}$ into the free group $\pi_1(F)$. Then $(f|_J)_*$ is multiplication by an integer $n \neq 0$.

If F has more than 1 boundary component, we can choose generators for $\pi_1(F)$ such that the generator of $\pi_1(J)$ goes to a generator of $\pi_1(F)$ under the left vertical map in (2). Since f_* is an isomorphism, it takes generators to generators, and thus it follows by commutativity that $n = \pm 1$.

If F only has the one boundary component J , then the generator α of $\pi_1(J)$ maps to either $\partial = \Pi_{i=1}^g[a_i, b_i] \in \pi_1(F)$ (oriented case) or $\partial = \Pi_{i=1}^g a_i^2 \in \pi_1(F)$ (unoriented case). If $f_*(\alpha) = x^n$ for a generator x of $\pi_1(K)$, we get by commutativity that $f_*(\partial) \in \pi_1(G)$ would be an n th power of something. Since $f_* : \pi_1(F) \longrightarrow \pi_1(G)$ is an isomorphism, ∂ itself would be an n th power of some element. In case $\partial = a_1^2$, F is a Möbius band, so this cannot happen. In all other cases we get $n = \pm 1$.

We have shown that $(f|_J)_* : \pi_1(J) \longrightarrow \pi_1(K)$ is an isomorphism. Thus we can homotope f in a collar neighborhood around J such that $f|_J : J \longrightarrow K$ is a diffeomorphism. We do this for every boundary component of F .

All that is left is to check that no two boundary components J_1, J_2 of F map to the same boundary component K in G . If that were the case, the elements of $\pi_1(F)$ generating $\pi_1(J_1)$ and $\pi_1(J_2)$ would both map to a generator of $\pi_1(K)$, i.e. would coincide up to a sign, since $f_* : \pi_1(F) \longrightarrow \pi_1(G)$ is an isomorphism. Then by Lemma 3.3, F would be a cylinder, which it is not. $\square$

Theorem 3.5. *Let F and G be compact surfaces, and let $f : F \longrightarrow G$ be a homotopy equivalence. Assume neither F nor G is a disk, a sphere, a cylinder, a Möbius band, a torus, a Klein bottle, or $\mathbb{R}P^2$. Then f is homotopic to a diffeomorphism.*

Proof. If F and G have non-empty boundary, Lemma 3.4 and Theorem 3.1 give the result. So assume that F and G are closed surfaces.

Let $B \subseteq G$ be a non-separating, 2-sided simple closed curve in G . Since f is homotopic to a smooth map which is transverse to B , we can assume that f is smooth and transverse to B . Consider the components of $f^{-1}(B)$. By transversality and compactness, this is a finite set of disjoint 1-submanifolds of F . As in the proof of Theorem 3.1, we can homotope f so that no component in $f^{-1}(B)$ bounds a disk. For any 1-sided simple closed curve γ in $f^{-1}(B)$, take a small tubular neighborhood M of γ such that $f(M) \subseteq N$, where N is a tubular neighborhood of B . Since $M \setminus \gamma$ is connected and $f(M \setminus \gamma) \subseteq N \setminus B$, it follows that $M \setminus \gamma$ maps to the same side of the 2-sided curve B under f . This implies that we can homotope f in M to a function not hitting B . So we can assume that no component of $f^{-1}(B)$ is a 1-sided simple closed curve.

Now let H_0, H_1 be two components of $f^{-1}(B)$, and let $h_0, h_1 : S^1 \rightarrow F$ be parametrizations of H_0 and H_1 , respectively. Then

$$\mathbb{Z} \cong \pi_1(H_i) \xrightarrow{f} \pi_1(B) \cong \mathbb{Z}$$

is multiplication by some $m_i \in \mathbb{Z}$. Note that $m_i \neq 0$ since h_i is nontrivial in F and f is injective on $\pi_1(F)$. This gives that $f_*(h_0^{m_1}) = f_*(h_1^{m_0}) \in \pi_1(G)$, and since f is injective on $\pi_1(F)$, $h_0^{m_1} = h_1^{m_0} \in \pi_1(F)$. Then by Lemma 3.3 they bound a cylinder (if h_0 and h_1 bound a cylinder then so do h_0 and h_1). This cylinder might contain components of $f^{-1}(B)$, but since there are finitely many such components, we can take a cylinder whose intersection with $f^{-1}(B)$ is precisely its ends, call them h_0 and h_1 again.

Now the cylinder gives a homotopy $c : S^1 \times I \rightarrow F$ from h_0 to h_1 , and thus $f \circ c : S^1 \times I \rightarrow G$ is a homotopy in G , with $f(c(S^1 \times]0, 1[)) \cap B = \emptyset$. Thus we get a continuous map $\widetilde{f \circ c} : S^1 \times I \rightarrow G \setminus B$ into the cut-up surface $G \setminus B$. This is a homotopy between non-zero powers of boundary components of $G \setminus B$. Now by Lemma 3.3, if these two boundary components are distinct, $G \setminus B$ would be a cylinder. But this is impossible, since G is neither a torus nor a Klein bottle. This implies that both ends of the cylinder is mapped to the same boundary component in $G \setminus B$, and thus we can change f by a homotopy to remove h_0 and h_1 from $f^{-1}(B)$ without changing $f^{-1}(B)$ otherwise. We can now assume that $f^{-1}(B)$ is a single closed curve, since $f^{-1}(B) = \emptyset$ implies that $f_* : \pi_1(F) \rightarrow \pi_1(G)$ factors through $\pi_1(G \setminus B)$ but $\pi_1(G \setminus B) \rightarrow \pi_1(G)$ is not surjective. We can finally see that the curve $f^{-1}(B)$ is non-separating by Lemma 2.6, since B is non-separating and $f_* : H_1(F) \rightarrow H_1(G)$ is a group homomorphism.

Consider $f| : N(f^{-1}(B)) \rightarrow N(B)$, where $N(-)$ denotes a tubular neigh-

neighborhood. Then, using a method as in the proof of Lemma 2.2 on $f^{-1}(B)$ and a bump function to extend to $N(f^{-1}(B))$, one can see that f is homotopic to a map g with $g^{-1}(B) = f^{-1}(B)$, such that $g|_{N(g^{-1}(B))}$ is a smooth covering map (the number of sheets will be the degree of $f : f^{-1}(B) \rightarrow B$). So now we assume that f is a smooth covering map on a neighborhood of $f^{-1}(B)$.

Since $f_* : \pi_1(F \setminus f^{-1}(B)) \rightarrow \pi_1(\widetilde{G \setminus B})$ is injective ([Lyndon-Schupp] prop 5.1), we can choose a covering $\rho : \widetilde{G \setminus B} \rightarrow G \setminus B$ and lift f as in the diagram,

$$\begin{array}{ccc} & & \widetilde{G \setminus B} \\ & \nearrow \tilde{f} & \downarrow \rho \\ F \setminus f^{-1}(B) & \xrightarrow{f} & G \setminus B \end{array} \quad (3)$$

such that $\tilde{f}_* : \pi_1(F \setminus f^{-1}(B)) \cong \pi_1(\widetilde{G \setminus B})$. Moreover, ρ is a finite-sheet covering, since f maps (a parametrization of) $f^{-1}(B)$ to a non-zero multiple of (a parametrization of) B , and the number of sheets is locally constant. So $\widetilde{G \setminus B}$ is compact.

Now in a neighborhood of the boundary of $F \setminus f^{-1}(B)$, $\tilde{f}$ is a covering map, and $\tilde{f}_*$ is an isomorphism on π_1 . So $\tilde{f}$ is an embedding on a neighborhood of the boundary. By Corollary 3.2 on $\tilde{f} : F \setminus f^{-1}(B) \rightarrow \widetilde{G \setminus B}$, $\tilde{f}$ is homotopic to a diffeomorphism, relative to a neighborhood of the boundary. Glue up this diffeomorphism to a map $g : F \rightarrow G$ which will be homotopic to f , and be both a homotopy equivalence and a smooth covering map. The last two imply that g is a diffeomorphism $F \rightarrow G$. $\square$

4 Injectivity

In this section we will prove that the map in (1) is injective, i.e. if a diffeomorphism is homotopic to the identity, it is smoothly isotopic to the identity.

Definition 4.1. Let $f, g : I \rightarrow F$ be smooth embeddings into a surface F . We say that f and g form an "eye" if the following is satisfied:

- (i) $f(I) \cup g(I)$ bounds a disk in F .
- (ii) $f|_{[0, \varepsilon[} = g|_{[0, \varepsilon[}$, $f|_{]1-\varepsilon, 1]} = g|_{]1-\varepsilon, 1]}$, and f is disjoint from g on $] \varepsilon, 1 - \varepsilon [$.

Lemma 4.2. Let $f, g : I \rightarrow F$ be two smooth embeddings into a surface F which form an "eye". Then there is a smooth isotopy φ_t of F with $\varphi_0 = \text{id}_F$ and $\varphi_1 \circ g = f$. Furthermore, there is a small neighborhood N of the disk bounded by f and g for which φ_t is the identity outside N for all t .

Proof. Let N_f be a tubular neighborhood of $f(I)$, given by a normal vector field ξ_f to N_f . Let also N_g be a tubular neighborhood of $g(I)$ given by a normal vector field ξ_g , such that $N_f \cup N_g$ is an annulus. This is possible since $f(I) \cup g(I)$ bounds a disk in F .

There is a diffeomorphism $\psi_f : N_f \longrightarrow V_f \subseteq \mathbb{R}^2$ such that $\psi_f \circ f$ is the standard embedding $I \longrightarrow \mathbb{R} \times \{0\}$. We can take $V_f = I \times]-\varepsilon, \varepsilon[$. We want to extend ψ_f to a diffeomorphism $\psi_{fg} : N_f \cup N_g \hookrightarrow \mathbb{R}^2$, i.e. $\psi_{fg}|_{N_f} = \psi_f$.

First we note that inside $V_f = I \times]-\varepsilon, \varepsilon[$ we have the image

$$G = \psi_f(g(I) \cap N_f).$$

By taking ε small, we can ensure that G is the graph $\{(t, h(t))\}$ of a smooth function $h : [0, \delta[\cup]1 - \delta, 1] \longrightarrow [0, \infty[$. We can extend ψ_f to a map $\tilde{\psi}_{fg}$ defined on $N_f \cup g(I)$ such that $\tilde{\psi}_{fg} \circ g : I \longrightarrow \mathbb{R}^2$ is smooth, using bump functions etc as usual, such that the image $G_I = \tilde{\psi}_{fg} \circ g(I)$ is the graph $\{(t, h(t))\}$ of a function $h : I \longrightarrow [0, \infty[$, see Figure 1.




Figure 1: The tubular neighborhood V_f and the graph G_I of h in $\mathbb{R}^2$.

We define a tubular neighborhood of G using the vector field $\eta_G = (\psi_f)_*(\xi_g|_{N_f \cap N_g})$. Since ψ_f is a diffeomorphism, η_G is a transverse vector field, and so defines a tubular neighborhood N_G of G inside V_f . Now we shrink V_f to $I \times]-\varepsilon', \varepsilon'[$ where $\varepsilon' < \varepsilon$ (thus also shrinking N_f). Then we cover G_I by two open sets in $\mathbb{R}^2$, U_1 covering $G_I \setminus G$, and U_2 whose intersection with U_1 lie in N_G and outside V_f , see Figure 2. Then we take a partition of unity ρ_1, ρ_2 with respect to U_1, U_2 .

Let η_I be the standard normal vector field to G_I , defined on $G_I \setminus G$. Then we make a new vector field $\rho_1 \eta_I + \rho_2 \eta_G$. Since $\rho_1 \eta_I + \rho_2 \eta_G$ is never 0 or tangent to G , this defines a tubular neighborhood V_g of G_I . This tubular neighborhood coincides with N_G on V_f , and thus gives a diffeomorphism $\psi_{fg} : N_f \cup N_g \longrightarrow V_f \cup V_g$ which extends $\tilde{\psi}_{fg}$.

The inner boundary circle C of the annulus $N_f \cup N_g$ bounds a disk $D' \subseteq F$, and so the image $\psi_{fg}(C)$ also bounds a disk $D_{\mathbb{R}^2} \subseteq \mathbb{R}^2$. Then we can extend




Figure 2: Neighborhoods U_1 and U_2 of G_I .

$\psi_{fg}|_C$ to a map $D' \rightarrow D_{\mathbb{R}^2}$, which is necessarily a homotopy equivalence, so by Lemma 2.4 we can replace it by a diffeomorphism $\psi_{D'} : D' \rightarrow D_{\mathbb{R}^2}$, such that $\psi_{D'}|_C = \psi_{fg}|_C$. Then we can glue $\psi_{D'}$ and ψ_{fg} along C to obtain a diffeomorphism Ψ from $D = D' \cup N_f \cup N_g$ onto a disk in $\mathbb{R}^2$.

Now we can use a vertical flow in $D' \cup N_g \cup N_f$ (i.e. a pullback under Ψ of the obvious vertical flow in $\mathbb{R}^2$) to make $\text{Im}(g) = \text{Im}(f)$, and lastly a horizontal flow in N_f to make $g = f$. $\square$

Lemma 4.3. *Given two smoothly embedded arcs $f, g : I \rightarrow F$ satisfying $f(\{0, 1\}) \cap g(I) = f(I) \cap g(\{0, 1\}) = \emptyset$. Then there is a smooth isotopy φ_t of $\text{id}|_F$ such that $\varphi_1 \circ f$ and g intersect transversally. Moreover φ_t is the identity outside a tubular neighborhood of f .*

Proof. Take an open tubular neighborhood of f , N_f , of constant radius, where $r : N_f \rightarrow f(I)$ is the retraction. Inside N_f take a closed tubular neighborhood of f of constant radius, N_f^c . We cover $g(I) \cap N_f^c$ with sets of the form $N_f(a, b) = \{x \in N_f \mid f^{-1}(r(x)) \in]a, b[\}$, where $a < b \in I$, and $f(a), f(b)$ is outside $g(I)$. Since $g(I) \cap N_f^c$ is compact, we can assume that it is a finite covering, $N_f(a_i, b_i)$, $i = 1, \dots, N$, where $a_1 < a_2 < \dots < a_N$. For each $x \in F$ where f and g intersect non-transversally, $x \in N_f(a_i, b_i)$ for some i . Now take the first such i . Then we can choose another arc $\tilde{g} : I \rightarrow N_f(a_i, b_i)$ such that g and $\tilde{g}$ form an "eye" and $\tilde{g}$ and f intersect transversally for all $x \in \tilde{g}(I) \cap f(I) \subseteq N_f(a_i, b_i)$. Now by Lemma 4.2 there is an isotopy from g to $\tilde{g}$ in $N_f(a_i, b_i)$, which is the identity outside $N_f(a_i, b_i)$. Doing this for each i , we obtain in finitely many steps an isotopy which is the identity outside N_f , making f and g intersect transversally. $\square$

Lemma 4.4. *Let F be a compact surface with $F \neq \mathbb{R}P^2, S^2$, and let f be a diffeomorphism of F .*

- (i) *Let $\alpha_i : S^1 \rightarrow F \setminus \partial F$ be a finite family of disjoint, non-trivial, pairwise non-homotopic two-sided simple closed curves, with $f \circ \alpha_i \simeq \alpha_i$ for all*

i. Then there is an smooth isotopy f_t of F such that $f_0 = f$ and $f_1 \circ \alpha_i = \alpha_i$ and the identity extends to tubular neighborhoods.

(ii) Let $\alpha_i : I \longrightarrow F$ be a finite family of simple curves, disjoint except possibly at endpoints, with $f \circ \alpha_i \simeq \alpha_i$ and $f \circ \alpha_i = \alpha_i$ near the endpoints for all i . Let $A \subseteq F$ be a union of disjoint non-trivial closed curves, with $f|_A = \text{id}$ and $\alpha_i(I) \cap A = \alpha_i(\partial I)$ for all i . Then there is an smooth isotopy f_t of F , such that $f_0 = f$, $f_1 \circ \alpha_i = \alpha_i$ and the identity extends to tubular neighborhoods. Furthermore $f_t|_A = \text{id}$ for all t .

Proof. (i) and (ii) can be proved by the same methods, so we handle the two cases as one initially. But we will also use (i) to prove (ii). First, in both cases we have a closed subset $A \subseteq F$ with $f|_A = \text{id}$ (in case (i), A starts as $\emptyset$). Consider a single curve $\alpha = \alpha_1$. We will make an isotopy f_t of F such that $f_0 = f$, $f_1 \circ \alpha = \alpha$, and $f_t|_A = \text{id}$ for all t . Then we can let $A_1 = A \cup \alpha(I)$, and use the result for f_1 and A_1 on α_2 , completing the proof in a finite number of steps. So consider a curve α as in (i) or (ii), and let $\beta = f \circ \alpha$ be the image curve. By assumption, $\beta \simeq \alpha$.

In case (ii), there are small neighborhoods N_0 and N_1 of the start and end points where α and β agree. Inside N_0 and N_1 we can make an isotopy of f which perturbs β slightly, so that α and β agree near the start/end point, and then become disjoint. By shrinking N_0 and N_1 we can assume that α and β are disjoint on ∂N_0 and ∂N_1 . Our goal is now to make α and β disjoint outside N_0 and N_1 . From now on, we will ignore N_0 and N_1 in the proof, and only work with α and β outside them.

By Lemma 4.3 we can assume α and β are transverse to each other. Then α and β have finitely many intersection points by compactness. To get an isotopy of F taking β to α , we will first ensure that α and β have no intersection points. To do this, consider the universal covering $\pi : \tilde{F} \longrightarrow F$. We can model $\tilde{F}$ as an open disk in $\mathbb{R}^2$. Take a fixed lift $\tilde{\beta}$ of β .

We consider all the connected components of $\pi^{-1}(\alpha)$ that intersect $\tilde{\beta}$. There are finitely many such components, call them $\tilde{\alpha}_k$, since α and β have finitely many intersection points. The $\tilde{\alpha}_k$ are also transverse to $\tilde{\beta}$. Now we look for a pair of intersection points between $\tilde{\beta}$ and an $\tilde{\alpha}_i$, such that the part of the two curves between these points (a closed curve, call it σ) bounds a disk whose interior does not contain any points on $\tilde{\beta}$ or $\tilde{\alpha}_k$ for any k . So σ is a simple closed curve in $\tilde{F}$ bounding a disk. Projecting onto F , we get $\pi \circ \sigma$ (the parts of α and of β between two intersection points) also a simple closed curve, which is null-homotopic, so according to Lemma 2.4, $\pi \circ \sigma$ bounds a disk in F . We can choose a curve β' which form an "eye" with β and which does not intersect α in a neighborhood of the disk bounded by $\pi \circ \sigma$. Then by lemma 4.2 we can isotope β to β' , so that there are two fewer intersection

points between α and β' . Since there are finitely many intersection points, this procedure terminates.

But we must show why we can always find such a σ in $\tilde{F}$. Since $\tilde{\alpha}_k$ is a connected component of $\pi^{-1}(\alpha)$, each $\tilde{\alpha}_k$ separates $\tilde{F}$. So if $\tilde{\beta}$ crosses $\tilde{\alpha}_i$ once, it must cross it again (let us choose the first time it does so), as it is transverse to $\tilde{\alpha}_i$. Now $\tilde{F} \subseteq \mathbb{R}^2$, so the part of $\tilde{\beta}$ and $\tilde{\alpha}_i$ between these two intersection points will bound a disk. If this disk contains parts of $\tilde{\beta}$ or $\tilde{\alpha}_k$'s, there will be a smaller disk inside which satisfies the requirements, since there are finitely many intersection points. In this way we can isotope β to a curve which does not intersect α (in case (ii), except in N_0 and N_1).

In case (i), we now have two homotopic disjoint simple closed curves α and β . Then according to Lemma 1.4, they bound a cylinder. Recall that the set A (fixed by f) consists of the curves already handled, i.e. a union of non-trivial closed curves, none of which are homotopic to α , and thus not to β , either. Thus A cannot intersect the cylinder bounded by α and β (in fact, A cannot intersect a small open neighborhood of the cylinder). Then clearly there is an isotopy f_t of F , which is the identity on A , taking β to α .

In case (ii), the two curves α and β are homotopic and form a simple closed curve, so again they bound a disk. Recall that A originally consisted of non-trivial closed curves, so none of these can be inside the disk. As we add curves to A , the circles get connected by arcs. None of these can intersect α , since they were assumed to be disjoint from the start. As f is the identity on A , they cannot intersect $\beta = f \circ \alpha$, either. Thus A cannot cross the boundary of the disk, so A and the disk are disjoint. Thus by lemma 4.2 we can make an isotopy f_t of F , which is the identity on A , so that $f_1 \circ \alpha = \alpha$.

Now we extend the result to tubular neighborhoods of the curves. We make a tubular neighborhood M_0 of α , and by compactness identify it with $S^1 \times]-\varepsilon, \varepsilon[$ in case (i) and $I \times]-\varepsilon, \varepsilon[$ in case (ii). Now for (x, t) in a smaller neighborhood $M_1 \subset M_0$ of α , the projection the second coordinate $\text{pr}_t f_x(t) := \text{pr}_t(f(x, t))$ has positive differential, and thus for all x the image of $f_x(t)$, $\{(x', t') \mid (x', t') = f_x(t) \text{ for some } t \in]-\varepsilon, \varepsilon[\}$ is the graph of a function $h_x(t') = x'$. Now we can make tubular neighborhood M_2 such that $M_2 \subset f(M_1)$ and by possibly shrinking it assume that $M_2 = I \times]-\delta, \delta[$ or $M_2 = S^1 \times]-\delta, \delta[$. For definiteness, say $M_2 = I \times]-\delta, \delta[$. Choose a smooth bump function $\rho(t)$ with $\rho(t) = 1$ for $|t| \leq \frac{1}{2}\delta$ and $\rho(t) = 0$ for $|t| = \delta$. Let

$$g_s(x, t) = \begin{cases} ((1-s)h_x(t') + s(\rho(t')x + (1-\rho(t'))h_x(t')), t') & \text{for } (x, t) \in f^{-1}(M_2) \\ f(x, t) & \text{otherwise.} \end{cases}$$

where t' is the second coordinate of $f(x, t)$ as above. Then g_s defines an isotopy from f to a function g_1 with the property that $g_1(x, t) = (x, t')$ for

$t' \in]-\frac{\delta}{2}, \frac{\delta}{2}[$. Now by stretching the parameter t' in each interval $\{x\} \times]-\delta, \delta[$, we can assume that f is the identity on a (smaller) neighborhood. $\square$

Corollary 4.5. *If we in addition to the requirements in lemma 4.4 require that f is the identity on ∂F , then the isotopy can be assumed also to be the identity on ∂F .*

Proof. All the steps in the proof can be done away from the boundary. $\square$

Theorem 4.6. *Let $F \neq S^2, \mathbb{R}P^2$ and let $f, g \in \text{Diff}(F, \partial F)$ be homotopic. Then f and g are smoothly isotopic.*

To prove this I use the following result from [Smale] without proof.

Theorem 4.7 (Smale). *Let $f \in \text{Diff}(D^2, \partial D^2)$. Then f is smoothly isotopic to the identity, and if f is the identity on the boundary then so is the isotopy.*

Proof of Theorem 4.6. If we prove that $f^{-1}g$ is smoothly isotopic to the identity, we will have a smooth isotopy from g to f . Thus we can restrict our attention to the case $f \simeq \text{id}$.

Choose a pair of pants/annular decomposition of the surface F , i.e. a collection of disjoint simple closed curves $\alpha_i : I \longrightarrow F, i = 1, \dots, n$, in F . By Lemma 4.4 (i), f is smoothly isotopic to a map g , which is the identity on a tubular neighborhood of the α_i . In each pair of pants P , choose two curves that cut P up into a disk (for each annulus, choose one curve). By Lemma 4.4 (ii), there is an isotopy of F , which is the identity on the α_i , from g to a map h fixing a tubular neighborhood of the two curves in each pair of pants. Then we can use Smale's Theorem 4.7 on each disk, getting an isotopy to the identity. $\square$

Corollary 4.8. *If we in addition to the requirements in theorem 4.6 require that f and g is the identity on ∂F , then the isotopy can be assumed also to be the identity on ∂F .*

Proof. This is done as in theorem 4.6, except that we use corollary 4.5 instead of lemma 4.4, and using the fact that the isotopy in theorem 4.7 can be chosen to be the identity on boundary. $\square$

5 Proof of the Main Theorem

As explained in the introduction, we will use a result of Epstein to prove the statement about $\text{Top}(F, \{\partial F\})$:

Theorem 5.1 (Epstein). *Let F a compact surface and let $f : F \longrightarrow F$ be a homeomorphism homotopic to the identity. Then f is isotopic to the identity.*

Proof. This is a part of [Epstein] Thm 6.4, which states exactly this result, but for maps preserving a basepoint. And clearly, by an isotopy we can assume that f preserves any given point x_0 , and then f will be homotopic to the identity through maps preserving x_0 . $\square$

Now we are ready to prove the bijections of the Main Theorem 1.1:

$$\pi_0(\text{Diff}(F, \{\partial F\})) \xrightarrow{\cong} \pi_0(\text{Top}(F, \{\partial F\})) \xrightarrow{\cong} \pi_0(\text{hAut}(F, \{\partial F\}))$$

Proof of Theorem 1.1. Suppose F is not a sphere, a disk, a cylinder, a Möbius band, a torus, a Klein bottle, or $\mathbb{R}P^2$. Consider the composite map from (1),

$$\pi_0(\text{Diff}(F, \{\partial F\})) \longrightarrow \pi_0(\text{hAut}(F, \{\partial F\})). \quad (4)$$

According to Theorem 3.5, the map is surjective, and by Theorem 4.6, it is injective. Now all that is left is to show that

$$\pi_0(\text{Top}(F, \{\partial F\})) \longrightarrow \pi_0(\text{hAut}(F, \{\partial F\}))$$

is injective. But that is Theorem 5.1. $\square$

We now deduce Theorem 1.2:

Proof of Theorem 1.2. Suppose F is not a sphere, a disk, a cylinder, a Möbius band, a torus, a Klein bottle, or $\mathbb{R}P^2$.

Similar to the proof of Theorem 1.1, we consider the composite

$$\pi_0(\text{Diff}(F, \partial F)) \longrightarrow \pi_0(\text{hAut}(F, \partial F)).$$

We can assume $\partial F \neq \emptyset$, otherwise this is the Main Theorem. By Cor. 3.2, it is surjective, and by Cor. 4.8 it is injective. To prove the result, it suffices to show that

$$\pi_0(\text{Diff}(F, \partial F)) \longrightarrow \pi_0(\text{Top}(F, \partial F)) \quad (5)$$

is surjective. Consider the following fibration,

$$\text{Diff}(F, \partial F) \longrightarrow \text{Diff}(F, \{\partial F\}) \longrightarrow \text{Diff}(\partial F). \quad (6)$$

Here, $\text{Diff}(\partial F)$ is a semidirect product $\Sigma_n \ltimes \text{Diff}(S^1)$, where Σ_n denotes the symmetric group of permutations of n elements, and n is the number of

boundary components of F . We have of course a similar fibration for Top . We use $\text{Diff}(S^1) \xrightarrow{\cong} \text{Top}(S^1)$, and this implies

$$\text{Diff}(\partial F) \xrightarrow{\cong} \text{Top}(\partial F). \quad (7)$$

Now apply the long exact sequence of homotopy groups for the fibration (6) and its counterpart for Top . Using (7) and the Main Theorem, we get by the 5-lemma that the map (5) is surjective.

Now assume F is oriented. We can write $\text{Diff}(F, \{\partial F\})$ as the disjoint union

$$\text{Diff}(F, \{\partial F\}) = \text{Diff}_+(F, \{\partial F\}) \sqcup \text{Diff}_-(F, \{\partial F\}),$$

where the latter denotes the orientation-reversing maps. Similarly for Top and hAut . Since the maps in the Main Theorem respect this disjoint union, we immediately get the second part of 1.2.

By the same argument we can deduce the last part of 1.2 from the first part. $\square$

References

- [Epstein] D. B. A. Epstein *Curves on 2-manifolds and isotopies*, Acta Mathematica 115 (1966), 83-107.
- [Hatcher] A. Hatcher, *Algebraic Topology*, Cambridge University press, 2002.
- [Hempel] J. Hempel, *3-manifolds*, Ann. of Math. Princeton University press, 2002.
- [Lyndon-Schupp] R. Lyndon and P. Schupp, *Combinatorial Group Theory*, Springer, 1977.
- [Smale] S. Smale, Diffeomorphisms of the 2-sphere, Proc. Amer. Math. Soc., 10 (1959), 621-626.